


# Welfare and productivity of sows and litters housed in farrowing pens compared to farrowing crates



Clara Singh

Supervisor: Professor Paul Hemsworth

## Contents

	<i>page</i>
Abstract .....	2
1. Introduction .....	2
2. Methods .....	5
4. Results.....	9
5. Discussion .....	13
6. Conclusion .....	16
7. Acknowledgements .....	17
References .....	17

Appendix 1: Sow raw data

Appendix 2: Piglet raw data

Appendix 3: Piglet raw data: Time budgets of behaviour

## Abstract

The use of confinement housing in the pig industry is a contentious animal welfare issue and is beginning to be phased out in many countries. However the use of sow crates during farrowing and lactation is still standard practice and alternatives that improve welfare without compromising productivity and ease of management are under development. A simple loose pen design was trialled alongside a conventional crate system for the period of lactation beginning three days post farrowing and measurements made relating to sow and piglet behaviour, welfare and productivity.

32 sows and their litters (a total of 343 piglets) were studied, with behavioural data recorded weekly over the three weeks of lactation. The two housing systems were compared for piglet survival and weight gain, sow and piglet skin injuries, suckling behaviour and piglet time budgets of behaviour. Frequency, duration and interval between each suckling bout were recorded as well as the frequencies of sow-piglet interaction, displacement at the udder and number of piglets missing each bout. Piglet time budgets of behaviour were observed using instantaneous scan sampling, recording investigatory and social piglet behaviour including close contact with sow and play behaviour.

No significant differences were found in piglet survival ( $P=0.20$ ) or weight gain ( $P=0.15$ ) between treatments. Sows in farrowing pens obtained significantly more skin injuries than those in crates ( $P=0.03$ ) but treatment had no effect on incidence of skin injury for piglets ( $P=0.08$ ). Suckling bout duration ( $P<0.01$ ) and frequency ( $P<0.01$ ) decreased with week while intervals between bouts increased ( $P<0.01$ ) but showed no difference between treatments. Overall, more sow-piglet interaction around suckling was observed in farrowing pens ( $P<0.01$ ), the frequency of which also decreased with week ( $<0.01$ ). Farrowing pens were also associated with more displacements ( $P=0.04$ ), more piglets missing bouts ( $P<0.01$ ) and fewer bouts terminated by the sow ( $P=0.05$ ). Piglet time budgets of behaviour revealed significantly more play behaviour ( $P=0.03$ ) and interaction with the sow ( $P=0.02$ ) occurring in farrowing pens and more manipulative behaviour ( $P=0.02$ ) and inactivity ( $P=0.02$ ) in farrowing crates.

In summary, farrowing pens appeared to provide more social and environmental stimulation for piglets, resulting in more activity and social interaction, which may be interpreted as an improvement to welfare. The increased incidence of sow skin injuries may indicate a lack of adaptation or experience of sows in the farrowing pen environment. Evaluation of productivity was inconclusive because of the limited sample size.

## 1. Introduction

### *Confinement housing for reproductive sows*

The use of confinement housing for reproductive sows in the pig industry is a contentious animal welfare issue. In recent years, public and consumer pressure has driven a shift away from the use of such systems, with many countries introducing legislation to ban or limit the time that sows spend in physically and behaviourally restrictive environments. The use of gestation stalls has been banned in several European countries with a European Union (EU) wide ban to be implemented in 2013 (Council of Europe 2001). In Australia, gestation stalls will be phased out by 2017 (PISC 2008). However the use of sow crates for farrowing and lactation is still standard practice in Australia and are used by 95% of pork producers in the major pig producing countries of the EU (Johnson & Marchant-Forde 2009). The conventional farrowing crate severely restricts the movement of sows and was designed partly for ease of management but primarily to minimise the incidence of piglet crushing (Edwards 2002), which remains the most significant cause of early piglet death in all housing systems (Marchant et al. 2000).

### *Behavioural restriction and sow and piglet welfare*

Welfare concerns associated with the use of farrowing crates are largely based on the physical and behavioural restriction of sows, which prevents expression of sows' natural behavioural repertoire. Pigs are naturally active and gregarious animals and in the wild will forage over large home ranges. Wild female pigs typically live most of their life in small groups made up of related individuals of several generations. Just prior to parturition, however, sows will leave their group to seek a safe and isolated area to build a nest for

farrowing and nursing their piglets before re-joining the group some days later (Gonyou 2001). Maternal behaviour evolved to favour piglet survival and maturation through provision warmth and nourishment, protection from predators and vertical social learning. Sow crates used during farrowing and lactation in intensive piggeries prevent sows from satisfying many maternal motivations by severely restricting their capacity to move around, interact with their piglets and the environment and stand and lie comfortably. Research has shown that sows restrained in crates direct less behaviour toward their piglets, vocalise to piglets less and are less responsive to piglet vocalisations (Cronin, Simpson & Hemsworth 1996). Some aspects of maternal behaviour may be considered unnecessary under human care (for example nest building, protection of piglets from predators and provision of warmth where heat lamps are provided) and probably for that reason have previously been ignored in the design of modern housing for intensively farmed farrowing and lactating sows. Despite this, domesticated sows have retained strong motivations to perform such behaviours and given the opportunity will display protective maternal behaviour, forming strong and lasting bonds with their offspring (Gustafsson et al. 1999), the prevention of which has implications for sow welfare (Dawkins 1988). It is also becoming apparent that piglets reared with confined sows may be deprived of some benefits relating to social learning and development (Oostindjer et al. 2010; Oostindjer et al. 2011a). These issues have prompted development of alternative farrowing and lactation housing designs that provide greater space allowance and increased behavioural opportunities for sows and piglets.

Housing during the lactation period must consider the physical and behavioural needs of both the lactating sow and her young piglets. These needs differ significantly and housing system design does involve some trade-off between the two (Baxter, Lawrence & Edwards 2011b). Increasing space allowance may improve sow welfare by enabling sows to turn around, lie down more comfortably and display more maternal behaviour but can simultaneously increase the risk of piglet crushing. Earlier comparative studies have shown increased rates of mortality in loose farrowing pens compared to crates (Barnett et al. 2001; Edwards & Fraser 1997). The survival of piglets during lactation is a major consideration both in terms of welfare and productivity; their primary needs being warmth, access to udder (enabling adequate milk intake) and protection from crushing. However increasing space allowance for sows may not necessarily compromise piglet survival. With improvement in pen designs, such as the incorporation of anti-crushing bars and improved flooring materials, some more recent studies have shown similar mortality rates to those achieved in crated housing (Cronin et al. 2000; Gu et al. 2011). It seems that although the risk of death by crushing by the sow tends to be greater in loose-housed systems, piglets reared in crates are at greater risk of death from other causes resulting in overall comparable rates of mortality (KilBride et al. 2012; Weber et al. 2009).

In addition to housing factors, there is large variation in maternal ability between sows due to factors such as age, parity, experience and genetics. These factors seem to interact with environmental factors to influence piglet survival and welfare, with particular behaviours only being expressed under certain conditions (Baxter et al. 2011). The increased space in loose pens enables sows to freely interact with their piglets and display more maternal behaviour, which may produce benefits for piglets. Research has suggested that in some situations, the demonstration of maternal behaviour (eg. nest building activity, response to piglet distress calls, nose contact with piglets during posture changes and restlessness when piglets are removed) is negatively correlated with risk of piglet crushing (Andersen, Berg & Bøe 2005). However mortality alone is a crude measure both of piglet welfare and productivity. By solely focusing on mortality some important factors related to piglet welfare, growth and social development may be overlooked, particularly over the longer term. There is some evidence that roomier pens are associated with longer periods of milk let down, less fighting among piglets and greater weight gain (Pedersen et al. 2011). Greater interaction between sows and piglets may also facilitate social learning, assisting piglets in the transition from milk to solid feed at weaning (Oostindjer et al. 2010).

### *Alternative lactation housing*

The long-term projection in the pig industry is movement toward totally confinement-free systems, which provide a social and physical environment that enables reproductive sows and their litters to fulfil their natural behavioural motivations. However, in abandoning the use of farrowing crates, it is essential that the benefits they provide in terms of preventing piglet crushing and ease and safety of management are not lost. Various new designs have been trialled eg. (Baxter 1991; Bøe 1993; Cronin, Simpson & Hemsworth 1996). These pen designs were developed for use during farrowing and the whole of the lactation period. However they are typically expensive to install, costly in terms of space and require increased labour and specific management skills (Baxter, Lawrence & Edwards 2011a). Over the first few days of piglet life it is essential that humans are able to safely access animals when necessary. It is also the period in which piglets are most vulnerable to crushing by the sow. Retaining the use crates for farrowing and those first critical days may present an opportunity to improve welfare during lactation without risking an increase in piglet losses. A system that restricts the use of farrowing crates to parturition and the first few days of piglet life but uses alternative housing for the majority of the lactation period may be a valuable intermediary setup before transitioning to a totally confinement-free system, allowing design and management factors to be further improved. Converting to any new farrowing/lactation housing system in industry will be costly and must be justified in terms of improvement to animal welfare without compromising productivity. There has previously been some discrepancy between experimental and industry obtained data on piglet survival in alternative systems (Baxter, Lawrence & Edwards 2012). The need for thorough assessment prior to implementation is therefore vital.

### *Comparison of two systems*

The aim of the present study was to assess the viability (in terms of welfare and production) of replacing the conventional farrowing crate with an alternative system (the farrowing pen) for the lactation period, beginning 3 days post farrowing in a commercial piggery. The farrowing pen being trialled had a slightly increased total floor area, standard heated creep area and sloping bars along two walls. The major differences between the two housing systems were the increased floor space in the farrowing pens and that sows were unrestrained and could freely walk and turn around.

Assessment of productivity is relatively straightforward and generally based on measurements of litter size, mortality rate, growth rate and sow farrowing rates. In terms of productivity, it was hypothesised that:

1. Piglet survival is unaffected by housing treatment.
2. Piglets housed in pens during lactation gain more weight, producing larger piglets at weaning than those housed in crates.

Evaluation of welfare is somewhat more complex and therefore it is important that multiple, animal-based parameters are used. Due to practicality and lack of consensus on welfare evaluation, it is common in industry to assess animal welfare indirectly, relying on resource and management factors. Animal-based measures can be more difficult and time-consuming to measure in a standardised way but do provide direct evidence of animals' coping success (Barnett & Hemsworth 2009). This approach is underpinned by the definition of welfare described by Broom (2008) who stated that: 'welfare of an individual is its state as regards its attempts to cope with its environment.' Using this definition, observation of abnormal physiological functioning or behaviour may indicate a lack of adaptation or coping success. With the use of physiological measurements of health/fitness complemented by observations of sow and piglet behaviour, welfare assessment during lactation attempts to determine whether the biological and behavioural needs of both the sow and her piglets are being met. Welfare assessment focused on presence of skin lesions, suckling behaviour, frequency of sow-piglet interaction and piglet investigative, social and play behaviour. These were considered to be relevant signs of both sow and piglet welfare.

Assessment of skin injuries has been used previously and can provide evidence of aggression as well as a lack of adaptation to confinement when sows are unable to coordinate their movements in order to avoid injury. For example, the incidence of skin abrasions was found to be higher in gestating sows housed in stalls as opposed to pens (Karlen et al. 2007).

Suckling behaviour is relevant to assessment of both productivity and welfare since adequate milk intake by piglets is vital. Previous studies have found a positive correlation between stability of teat order and growth rate of piglets (Hemsworth, Winfield & Mullaney 1976). Improved access to the udder (as occurs in farrowing pens) has also been associated with increased weight gain and longer periods of milk let-down (Pedersen et al. 2011). Frequency of aggression and interruptions during nursing may also contribute to less effective milk intake by piglets and therefore piglet welfare.

Play behaviour can be used as an indicator of positive welfare (Boissy et al. 2007; Newberry, Wood-Gush & Hall 1988) and may be affected by housing conditions (Chaloupková et al. 2007). Welfare assessment typically focuses on provision of certain requirements and absence of signs of suffering. However the presence of positive indicators of welfare (such as play) are increasingly being recognised as important (Held & Špinka 2011).

By assessment of these welfare indicators, the present study aimed to demonstrate that loose-housing of sows in farrowing pens during lactation enhances welfare for both sows and piglets not only by physical means but by promoting interaction and bonding between sows and piglets.

Relating to welfare, the following hypotheses were tested:

1. Farrowing pen housing during lactation is associated with fewer skin lesions than in farrowing crates for both sows and piglets.
2. Improved social contact and udder access in farrowing pens is associated with more effective nursing with longer suckling bouts, fewer displacements during suckling and fewer piglets missing suckling bouts.
3. Increased area and freedom of movement in farrowing pens provides increased opportunity for sows and piglets to interact during and around suckling bouts resulting in more frequent sow-piglet nose contact.
4. Increased area and freedom for social interaction in farrowing pens results in more frequent investigative, social and play behaviours and less frequent manipulative and aggressive behaviour in piglets.

## **2. Methods**

All animal procedures were conducted with prior approval obtained through The University of Melbourne Animal Ethics Committee.

### *2.1 Animals and husbandry*

Productivity and skin injury assessments were made on 32 sows of mixed parity and their litters (a total of 343 piglets). Each sow and litter unit was randomly allocated to either a control (farrowing crates) or experimental (farrowing pens) treatment group. Twenty-four of the sows and 96 piglets (4 per litter) were observed for behavioural analysis. Focal piglets were selected randomly, balancing for sex and excluding any obvious runts. The study took place at Berrybank Farm, Ballarat, Victoria where all aspects of daily animal husbandry were managed by farm staff according to their normal practice.


### *2.2 Experimental design*

The experiment was run as four identical time replicates (due to small number of available pens) with each replicate consisting of eight sows (four per treatment). All sows were housed in groups during gestation and moved to farrowing crates for parturition. The experimental period began at three days post partum (piglets approximately 3 days of age), when control and experimental sows and their litters were transferred to their respective housing treatments in the lactation shed. They remained there for a further 22-25 days until the piglets were weaned at around four weeks of age.

### *2.3 Housing treatments*

Control sows and their litters were housed in farrowing crates and experimental sows and their litters were housed in farrowing pens in the same shed (see figure 1). The crates and pens were located on the same side of the shed on opposite sides of the aisle. The farrowing crates used were typical of conventional systems, with a space allowance of approximately 1.2m<sup>2</sup> that did not allow sows to turn around. The lactation pens used for the experimental group provided a floor space of over 4m<sup>2</sup>. Both physiological and behavioural

data relating to productivity and welfare of sows and litters were obtained weekly over the period of lactation.



**Figure 1 - Farrowing crate (left) and farrowing pen (right) design**

#### **2.4 Measurement of productivity**

The following production data were recorded:


- litter size (ie. no. of piglets alive at beginning of experimental period)
- piglet weight at start of experiment/lactation (approx. 3 days of age)
- piglet weight at weaning
- number of piglet deaths during experiment/lactation

#### **2.5 Measurement of welfare**

Data relating to welfare were collected three times for each sow and litter at weekly intervals and consisted of sow and piglet skin injury assessments as well as behavioural observations from video footage.

##### **2.5.1 Skin injuries**

Sow and piglet skin injuries were scored using the technique described by Karlen et al. (2007). Only fresh lesions were recorded and these were defined as any injury that appeared red or inflamed indicating it had been obtained recently. Each lesion was described as either a scratch, abrasion, ulcer or cut and further categorised according to location on the body (see figure 2). Injury scores were recorded in the second and third weeks (therefore recording injuries obtained during the first and second weeks) of the lactation period with two assessments made for each sow and focal piglet.



**Figure 2 - Classification of body regions for skin injury assessment as used by (Karlen et al. 2007)**

### 2.5.2 Behaviour

Sow and piglet behaviour during lactation was recorded using both digital video recorders installed above each pen and a hand-held camcorder. Overhead cameras recorded continuously on the 2nd, 9th and 16th day of the experimental period between the daylight hours of 0700 and 1700 h. This footage was used for quantification of suckling behaviours and sow-piglet interactions around suckling bouts.

Hand-held recordings were obtained on the same days for four hours each day using a scan sampling technique in which pens/crates were filmed for 30s at a time, constantly rotating. Recordings were made for one hour at a time with hours starting at 0800, 0910, 1020 and 1130 h, resulting in a total of 60 30s scans per day per pen/crate. These recordings were used to obtain data on piglet behavioural time budgets.

All video footage was viewed retrospectively to quantify the following specific sow and piglet behaviours:

### 2.5.3 Behaviour: suckling behaviour

The following data was recorded for each suckling bout between 0700 and 1700 h on the 2<sup>nd</sup>, 9<sup>th</sup> and 16<sup>th</sup> day in lactation housing:

- Length of suckling bout
- Termination of suckling: by sow/piglets
- Number of displacements during suckling
- Number of piglets that missed milk let-down
- Number of sow-piglet or piglet-sow (recorded together) interactions for two minutes prior, during and two minutes after suckling bouts.

Table 1 below lists the definitions used to record the above listed parameters.

**Table 1 - Definitions used for quantification of suckling behaviour observed by video footage.**

Suckling bout	A bout of suckling where at least 75% gather at udder and massage/suckle for >1.5min. NB. 75% of piglets was calculated as 7/9, 8/10 or 9/11, depending on number of piglets in litter.
Start of suckling bout	75% of piglets gather at udder and begin massaging.
End of suckling bout	75% of piglets have stopped massaging (became still or moved away from the udder) OR sow terminates massage/suckling by changing posture.
Sow-piglet interaction	Close contact <10cm between sow's nose and piglet body, where sow moves her head to make contact with piglet, pausing and maintaining close contact for >1s.  Or Close contact <10cm between piglet's nose and sow's face where piglet moves to make contact, pausing and maintaining close contact for >1s. Interactions were recorded as single events when close contact was observed repeatedly within a period of 5s between the same piglet and sow
Piglets that missed suckling bout	Piglets absent from udder (not suckling/massaging or attempting to) for a majority of suckling bout duration.
Displacement at udder	A piglet is displaced from the udder (becomes detached from the teat and has to walk outside or over the cluster of piglets to claim new position) during suckling bout, when 75% piglets are still suckling/massaging and prior to milk letdown (75% piglets still, finished when first piglets resumes massage). Does not include piglets that leave the udder and do not return immediately and resume suckling/massage

#### 2.5.4 Behaviour: piglet time budgets of behaviour

Activity of each of the four focal piglets from each litter was recorded for each 30s interval to estimate frequencies of significant behaviours as described in Table 2.

**Table 2 - Definitions of piglet behaviours used to estimate piglet time budgets of behaviour, adapted from the definitions used by Oostindjer et al. (2011c).**

Lying/inactive	Lying or sitting, not engaged in activity.
Walking/standing	Standing or walking but not otherwise engaged in activity (includes urinating and defecating).
Suckling/udder massage	Either mouth on teat or nose contact to udder with vertical head movements.
<i>Investigatory behaviours</i>	
- Investigating pen/floor/sow body	Sniffing, touching, biting, rooting, pawing or rubbing any part of the pen, floor or sow's body other than head.

– Investigating food/water	Sniffing, touching, pawing, eating or drinking food or water.
<i>Social behaviours</i>	
– Nosing other piglet	Touching or sniffing any part of the head or nose of another piglet (distance <10cm).
– Manipulative behaviour	Nibbling, sucking, chewing or rubbing with nose any part of the body of another piglet.
– Aggression	Ramming, pushing or biting another piglet.
– Mounting	Standing on the back of another piglet with front legs.
– Play	Shaking head, pivoting, jumping, or running with bouncy or jerky movements.
– Interacting with sow	Sniffing, grunting or nuzzling sow’s head with <10cm between piglet nose and sow’s head.

## Data analysis

All data were analysed using the statistical package, SPSS Statistics. Analysis of variance for repeated measures was used to examine between groups (treatment) effects as well as within subject (time) effects on sow and piglet behavioural measures and injury scores. Piglet survival and weight gain data were analysed using univariate analysis of variance. All data were tested for normality prior to analysis and a square root transformation was applied to all data sets that were non-normally distributed.

## 4. Results

### 4.1 Measurement of productivity (Table 3)

Analyses of piglet weight gain and survival were conducted on results from individual piglets and litters rather than overall means. Piglet daily weight gain and survival were lower for animals housed in farrowing pens compared to crates, however these differences were not significant (P=0.15 and P=0.20 respectively).

**Table 3 - Effects of housing treatment on piglet daily weight gain and survival during lactation**

Main effects	Means		SEM	F <sub>1,117</sub>	P value
	Crates	Pens			
Piglet % daily weight gain*	3.57 (12.74)	3.33 (11.09)	0.16	2.12	0.15
				F <sub>1,30</sub>	
Piglet % survival/litter*	9.71 (94.24)	9.55 (91.17)	0.12	1.74	0.20

\*Data square root transformed prior to statistical analysis. Back-transformed means are presented in parentheses.

### 4.2 Measurement of welfare

#### 4.2.1 Skin injuries (Table 4)

Skin injury analysis was conducted using results from individual sows and focal piglets. The incidence of skin injury for sows was significantly higher in pens compared to crates (P=0.03). There was no effect of week (P=0.49) or any effect of a week × treatment interaction (P=0.95). Housing treatment had no effect on the incidence of piglet skin injury (P=0.08) however there were significant effects of both week (P<0.01) and an interaction effect between week and treatment (P=0.01). Piglets had significantly more injuries in the third week of lactation compared to the second and this effect was more pronounced in pens than in crates.

The majority of both sow and piglet skin injuries obtained during the lactation period were minor scratches (sows mean 73% scratches, 21% abrasions and piglets mean 97% scratches). In the piglets, most of these injuries (86%) were on the ears, face, shoulders and neck regions (regions 1, 2, 3, 4 and 5). Sows showed a similar pattern of injuries, with a large proportion (30%) occurring around the face, neck and ears (regions 1, 2, 3 and 4). Neither sows nor piglets in either housing treatment obtained serious injury during the experimental period.

**Table 4 - Effects of housing treatment on the incidence of sow and piglet skin injury during the lactation period**

Main effects	Week	Means		SEM	F <sub>1,30</sub>	P value
		Crates	Pens			
Total new injuries (sows)*	2	0.53 (0.28)	1.11 (1.23)	0.32	5.05	0.03
	3	0.65 (0.43)	1.25 (1.57)	0.34		
F <sub>1,124</sub>						
Total new injuries (piglets)*	2	0.51 (0.26)	0.46 (0.21)	0.14	3.06	0.08
	3	1.62 (2.63)	2.19 (4.79)	0.23		

\*Data square root transformed prior to statistical analysis. Back-transformed means are presented in parentheses.

#### 4.2.2 Suckling behaviour (Table 5)

Suckling behavioural data were analysed as means per sow/litter unit. Neither the number of suckling bouts, duration of suckling nor duration of the interval between bouts differed significantly between treatments. Week had a significant effect on both the number of suckling bouts (P<0.01) and suckling bout duration (P<0.01), with the average number of bouts highest in the second week and decreasing in the third week. Accordingly, the duration of intervals between suckling bouts also decreased in the second week and increased again in the third week for both treatments (P<0.01). Suckling bout duration declined over time in both housing treatments. No week × treatment interactions were evident for these suckling behaviour variables.

Frequency of sow-piglet interactions occurring during the two minutes immediately prior to suckling was unaffected by housing treatment (P=0.38), week (P=0.93) or a week × treatment interaction (P=0.06). However, interactions during suckling bouts did show a treatment effect with a greater number of interactions observed in farrowing pens (P<0.01). Week also showed an effect (P<0.01), with the frequency of interactions decreasing over time. There was no interaction effect between week and treatment (P=0.22). Interactions observed in the two minutes immediately post suckling showed similar effects; with significantly more frequent interaction in the farrowing pens (P=0.01). There was also a significant interaction effect between week and treatment (P<0.01) but no overall effect of week (P=0.88). The number of interactions observed post suckling in farrowing pens increased with time while those in farrowing crates decreased. When total interactions around suckling were analysed as a single measure (as a sum of pre, during and post suckling interactions), frequencies generally declined over time in both pens and crates (P<0.01) but much more substantially in the farrowing crates, showing a significant interaction effect

between treatment and week ( $P<0.01$ ). There was also significantly more interaction around suckling occurring in the pens than the crates overall ( $P<0.01$ ).

The number of piglets that missed suckling bouts was higher in farrowing pens than crates ( $P<0.01$ ). This variable also showed a week ( $P=0.02$ ) and a week  $\times$  treatment interaction ( $P=0.04$ ) effect. The average number of piglets missing bouts generally decreased over time in both treatments but the effect was more pronounced in the crates. Similar effects were seen in the frequency of displacements. Overall, displacements were more frequent in the pens ( $P=0.04$ ) and declined over time in both treatments ( $P<0.01$ ), also showing a week  $\times$  treatment effect ( $P=0.05$ ), decreasing more rapidly in farrowing pens.

The proportion of bouts terminated by the sow was significantly affected by both housing treatment ( $P=0.05$ ) and week ( $P=0.03$ ). Sows housed in crates terminated a greater proportion of suckling bouts and in both treatments there was an overall increase in that proportion over time apart from a marked reduction in week two in the pen treatment. No week  $\times$  treatment effect was evident ( $P=0.76$ ).

**Table 5 - Effects of housing treatment on sow and piglet suckling behaviour**

Main effects	Week	Means		SEM	F <sub>1,22</sub>	P value
		Crates	Pens			
No. of suckling bouts	1	14.00	14.92	0.95	2.28	0.15
	2	14.67	15.83	0.93		
	3	12.75	13.83	0.79		
Bout duration (s)*	1	21.48 (461.36)	20.19 (407.53)	1.19	0.80	0.38
	2	19.91 (396.36)	19.63 (385.49)	0.71		
	3	17.98 (323.45)	17.66 (311.71)	0.84		
Bout interval (s)	1	2636	2438	198	2.55	0.13
	2	2488	2299	189		
	3	2867	2654	213		
Sow-piglet interactions 2min pre-suckling*	1	1.27 (1.61)	1.18 (1.40)	0.15	0.80	0.38
	2	1.09 (1.18)	1.30 (1.70)	0.13		
	3	1.05 (1.10)	1.36 (1.86)	0.12		
Sow-piglet interactions during suckling*	1	1.78 (3.17)	1.99 (3.94)	0.20	10.17	<0.01
	2	1.32 (1.74)	1.88 (3.55)	0.15		
	3	1.19 (1.41)	1.17 (1.36)	0.14		
Sow-piglet interactions 2min post-suckling	1	1.77	1.63	0.24	7.90	0.01
	2	1.43	1.89	0.27		
	3	1.09	2.42	0.35		
TOTAL sow-piglet interactions around suckling*	1	2.59 (6.69)	2.68 (7.17)	0.18	13.83	<0.01
	2	2.11 (4.46)	2.69 (7.22)	0.14		
	3	1.92 (3.68)	2.64 (6.96)	0.14		
Piglets missed suckling bouts*	1	1.92 (3.68)	2.64 (6.96)	0.08	12.55	<0.01
	2	0.44 (0.20)	0.47 (0.22)	0.08		

Main effects	Week	Means		SEM	F <sub>1,22</sub>	P value
		Crates	Pens			
	3	0.22 (0.05)	0.50 (0.25)	0.09		
Displacements during suckling*	1	1.16 1.36	1.63 (2.65)	0.26	4.79	0.04
	2	0.64 (0.41)	1.05 (1.11)	0.16		
	3	0.77 (0.59)	0.68 (0.46)	0.12		
Proportion of bouts terminated by sow*	1	0.68 (0.47)	0.60 (0.36)	0.07	4.49	0.05
	2	0.68 (0.47)	0.55 (0.30)	0.06		
	3	0.77 (0.59)	0.67 (0.45)	0.07		

\*Data square root transformed prior to statistical analysis. Back-transformed means are presented in parentheses.

#### 4.2.3 Piglet time budgets of behaviour (Table 6)

Frequencies of piglet behaviours were expressed as proportions, that is, number of times each behaviour was observed divided by total number of observations and the means are given in Table 6 below. Data analyses were conducted on results from individual piglets. Treatment effects were apparent in the frequencies of several piglet behaviours; lying (P=0.02) and manipulative behaviours (P=0.02) were more frequently observed in farrowing crates whereas play behaviour (P=0.03) and close contact with sow (P=0.02) were observed more frequently in farrowing pens. Many piglet behaviours were also affected by week with time spent lying (P<0.01) and suckling/massaging udder (P<0.01) showing a general reduction over time, although in farrowing pens the maximum mean value for proportion of time spent at the udder was calculated for the second week of lactation. Investigatory behaviours of investigating pen (P<0.01) and investigating/eating food (P<0.01) as well as social behaviours of nosing (P=0.03), aggression (P<0.01), manipulative behaviour (P<0.01) and play (P<0.01) all increased over the three observed weeks however investigating/eating food and nosing pen-mates were observed most frequently in the second week. Interaction effects between treatment and week were not apparent for most piglet behaviours, with lying behaviour showing an interaction effect (P=0.02) where lying decreased much more rapidly over time in the pens compared to crates.

**Table 6 - Effects of housing treatment on frequencies of piglet behaviours between 0800 and 1230 h over three weeks of lactation**

Main effects	Week	Means		SEM	F <sub>1,124</sub>	P value
		Crates	Pens			
Lying/inactive*	1	0.79 (0.62)	0.79 (0.63)	0.01	5.78	0.02
	2	0.74 (0.55)	0.72 (0.52)	0.01		
	3	0.75 (0.56)	0.71 (0.51)	0.01		
Walking/standing*	1	0.25 (0.06)	0.24 (0.06)	0.01	0.02	0.89
	2	0.26 (0.07)	0.25 (0.06)	0.02		
	3	0.23 (0.05)	0.24 (0.06)	0.02		
Suckling/udder massage*	1	0.49 (0.24)	0.48 (0.23)	0.02	0.01	0.90
	2	0.48 (0.23)	0.51 (0.26)	0.01		

Main effects	Week	Means		SEM	F <sub>1,124</sub>	P value
		Crates	Pens			
	3	0.46 (0.21)	0.46 (0.21)	0.02		
Investigating pen/floor/sow body*	1	0.17 (0.03)	0.16 (0.02)	0.02	0.22	0.64
	2	0.21 (0.04)	0.22 (0.05)	0.02		
	3	0.23 (0.05)	0.24 (0.06)	0.02		
Investigating food/water*	1	0.01 (0.00)	0.01 (0.00)	0.01	0.01	0.27
	2	0.57 (0.33)	0.27 (0.07)	0.01		
	3	0.11 (0.01)	0.11 (0.01)	0.02		
Nosing other piglet*	1	0.11 (0.01)	0.10 (0.01)	0.02	2.98	0.09
	2	0.13 (0.02)	0.15 (0.02)	0.02		
	3	0.11 (0.01)	0.14 (0.02)	0.02		
Manipulative behaviour*	1	0.02 (0.00)	0.01 (0.00)	0.01	5.68	0.02
	2	0.08 (0.01)	0.05 (0.00)	0.02		
	3	0.06 (0.00)	0.05 (0.00)	0.02		
Aggression*	1	0.01 (0.00)	0.01 (0.00)	0.01	0.91	0.34
	2	0.13 (0.02)	0.11 (0.01)	0.02		
	3	0.13 (0.02)	0.18 (0.03)	0.02		
Mounting*	1	0.00 (0.00)	0.00 (0.00)	0.00	0.30	0.58
	2	0.00 (0.00)	0.00 (0.00)	0.01		
	3	0.00 (0.00)	0.01 (0.00)	0.01		
Play*	1	0.02 (0.00)	0.02 (0.00)	0.01	4.97	0.03
	2	0.05 (0.00)	0.07 (0.00)	0.02		
	3	0.04 (0.00)	0.07 (0.01)	0.02		
Interacting with sow*	1	0.05 (0.00)	0.08 (0.01)	0.02	6.174	0.02
	2	0.08 (0.01)	0.09 (0.01)	0.02		
	3	0.08 (0.01)	0.11 (0.01)	0.02		

\*Data square root transformed prior to statistical analysis. Back-transformed means are presented in parentheses.

## 5. Discussion

### 5.1 Productivity

No significant differences in piglet survival or growth were apparent in the present study, although mean piglet daily weight gain and survival were slightly greater in animals housed in farrowing crates. These results support the hypothesis that survival is similar in the two treatments but do not support the hypothesis that growth rate is increased in farrowing pens. However, larger numbers of litters would be necessary to rigorously test these two hypotheses, particularly to test for differences in rates of survival. A tendency for slightly increased mortality rates for piglets reared in farrowing pens has been demonstrated in previous research comparing crated and loose-housed sow systems eg. (Blackshaw et al. 1994; de Oliveira Junior et al. 2011; Glastonbury 1976). Increased space for sows has been associated with greater risk of overlay and chilling of piglets (Edwards & Fraser 1997). However, most research into alternative housing uses sows that have had no or limited prior experience in the new system being trialled. Sow experience could potentially make a significant difference to piglet survival and welfare in alternative systems and this effect has been observed in the trial of a group farrowing system (Wechsler 1996). In the present study it was not possible to control or balance for sows who had reared litters in farrowing pens previously.

Further research is required to examine the effects of both farrowing pen housing and sow experience on sow productivity. Productivity evaluation of large numbers of animals is essential prior to implementation of any new system as even marginal differences in productivity can have a highly significant economic impact on the farm.

## *5.2 Sow welfare*

### *5.2.1 Injuries*

The interpretation of sow skin injury scores with regard to sow welfare requires some knowledge as to the cause and severity of those injuries. When sows are housed individually, skin injuries may be obtained by scrapes or bumps on pen/crate walls or bars and/or from contact with piglets. In the present study, the majority of the sow injuries obtained were minor scratches and abrasions. The difference in incidence of skin injury between housing treatments may reflect a lack of experience of sows changing posture in the farrowing pens, particularly when changing from standing to lying. With a large proportion of injuries present around the sows' faces and necks it is also plausible that the increased interaction with piglets that occurred in the farrowing pens contributed to more scratches, as piglets were observed climbing on, playing with and biting sows, particularly around their ears and face. Although the observed difference in sow skin injuries did not support the hypothesis that injuries would be reduced in farrowing pens, it may not indicate any significant negative impact on the welfare of sows housed in farrowing pens, since no serious injury was observed.

### *5.2.2 Expression of maternal behaviour*

Domestic sows retain many of the behavioural motivations of their wild ancestors and this is particularly apparent during farrowing and lactation. At this time the sow is highly motivated to perform specific behaviours relating to the care and protection of her offspring including nest-building, farrowing and nursing. Greater expression of maternal behaviour can be associated with improved piglet survival and weight gain but also depends on the environment (Andersen, Berg & Bøe 2005; Arey & Sancha 1996). Sow-piglet interaction seems to be an important expression of maternal behaviour during lactation and is necessary both for recognition of piglets by the sow and to coordinate effective suckling (Algers 1993). Sows mainly rely on olfactory cues to recognise their piglets and will often behave quite aggressively to unfamiliar piglets (Horrell 1982). In the present study, the increased interaction observed between sows and piglets in the farrowing pens (both around suckling and also overall as measured by piglet time budgets of behaviour) provides evidence of behavioural restriction of sows in crates. The frequency of sow-piglet interaction around suckling also decreased less over time and the overall frequency of piglet interactions with the sow increased more over time in the farrowing pens compared to crates, which may be a result of the formation of stronger social bonds between sows and piglets. Sow-piglet nasal contact has also been shown to occur more frequently in wild-type compared to domestic sows and piglets (Gustafsson et al. 1999) suggesting that it is a natural behaviour with some evolutionary advantage.

Maternal behaviour may have been limited in this study by the use of farrowing crates for parturition and the first three days of piglet life. Maternal behaviour is regulated by the release of specific hormones, the functioning of which involves complex mechanisms (Algers & Uvnäs-Moberg 2007) that could be interrupted by behavioural restriction. The prevention of early maternal behavioural expression, particularly nest-building, may have suppressed subsequent maternal behavioural motivation and expression in sows. Several studies have shown that suckling behaviour establishes more easily in litters in which the sow farrowed in pens with straw compared to standard crates (Cronin & Smith 1992). A correlation has also been reported between the expression of nest building behaviour and a reduced incidence of piglet crushing (Andersen, Berg & Bøe 2005; Pedersen et al. 2006). The importance of early maternal behavioural expression on subsequent behaviour requires further research.

The finding that sows in farrowing crates terminated more suckling bouts may indicate a less comfortable environment although this difference did not seem to affect the amount of time piglets spent at the udder, as no difference in bout duration was seen between treatments. It has been shown previously that sows tend to terminate more suckling bouts as piglets mature (Gustafsson et al. 1999) and this effect was seen in both treatments. The finding that sows in farrowing crates terminated more bouts could be another indication of sows' limited ability to communicate with their piglets caused by the physical restriction imposed by crates.

### *5.3 Piglet welfare*

#### *5.3.1 Piglet injuries*

The lack of treatment effect observed for piglet skin injuries did not support the hypothesis that fewer injuries would be obtained in farrowing pen housing. A majority of piglet injuries were minor scratches and the pattern of injury suggests most were obtained from fighting among pen-mates. This assumption fits with the finding that frequency of aggression also did not differ between treatments.

#### *5.3.2 Suckling behaviour*

The greater amount of sow-piglet interaction occurring around suckling bouts in farrowing pens supported the hypothesis that sows and piglets would interact by close nasal contact more frequently in farrowing pens. Increased interaction benefits piglets by facilitating effective nursing and may also aid piglets through vertical social learning. In a favourable environment, piglets will learn by observation and participation with the sow. For example, piglets can learn to eat solid food from the sow, assisting in the transition to solid feed at weaning (Oostindjer et al. 2011a). Close nasal contact between sows and piglets is thought to be important for recognition and communication and nursing behaviour has been shown to be regulated by pheromones and olfactory cues detected by piglets (Morrow-Tesch & McGlone 1990a; Morrow-Tesch & McGlone 1990b). The decline in frequency of nose-nose contact between sows and piglets around suckling over time was also expected and fits with the parent-offspring conflict theory, in which weaning occurs gradually as the sow invests less in caring for her young (Bøe 1993). Accordingly, suckling bout frequency and duration also decreased over time. The fact that nose contact with sows was more frequent in farrowing pens compared to crates can be explained as follows. Firstly, farrowing pens are less physically restrictive due to greater floor space and lack of crate bars, making the sow's head more accessible to piglets. Secondly, the loose-housing of sows may be more conducive to the expression of maternal behaviour by sows resulting in increased sow initiated close contact with piglets.

Contrary to what was hypothesised however, the number of displacements occurring during suckling and the number of piglets missing suckling bouts were greater in farrowing pens than crates. It is not clear why this may have been the case. The greater floor area in the farrowing pens might have meant that piglets were often sleeping at a greater distance from the sow and did not hear her grunting to initiate suckling. Similarly, the increased area may have allowed for more movement at the udder whereas in the crates, piglets had so little space it was not always possible to change teat position after initiation of suckling/massage.

#### *5.3.3 Piglet behaviour*

Significant differences in several parameters of piglet behaviour highlight the relevance of lactation housing to piglet behavioural expression and therefore welfare. The increased play

behaviour observed in the farrowing pens was likely a result of a combination of factors. The increased space may have made play behaviour physically more easily expressed. Increased space may have also provided more environmental and social stimulation for piglets, improving their overall welfare and therefore inclination to engage in play behaviour. It was interesting to note that although there was a significant difference in play behaviour between treatments, there was no difference in the frequency of aggression, which appeared to be mostly play-fighting. It is generally accepted that presence of play behavior is a strong indicator of positive welfare and these observed differences supported the hypothesis that increased area and freedom in farrowing pens is associated with increased play behaviour. However, the hypotheses that housing treatment would affect the frequency of nasal contact and aggression between piglets was not supported. The benefits of improved welfare during lactation may well persist beyond weaning. The effect of loose-housing of sows during lactation on piglet play behaviour seems to continue post weaning, with increased play behaviour expressed by piglets reared in loose-housing systems (Chaloupková et al. 2007; Oostindjer et al. 2011b).

In contrast to the effect the farrowing pen treatment had on play behaviour, manipulative behaviour was reduced and this effect has been reported previously when comparing loose and confined housing of sows. Oostindjer et al. (2011b) observed that lower levels of belly-nosing and other manipulative behaviours were associated with loose-housing of sows. Manipulative-type behaviour is thought to be redirected behaviour, occurring when the appropriate stimulus is inaccessible. Belly-nosing is one common manipulative-type behaviour often seen in early weaned piglets. It is an abnormal behaviour with no obvious function and is sometimes considered a stereotypy. High levels of such behaviour become a problem when skin injury and ulceration result and piglets that express high levels of belly-nosing have also been shown to have a reduced growth rate (Straw & Bartlett 2001). The increased manipulative behaviour observed in crate-housed piglets may indicate a lack of environmental enrichment and/or overcrowding (Dybkjær 1992). Similar to the findings of Oostindjer et al. (2011b), housing treatment did not appear to affect the frequency of nosing pen-mates, mounting or aggression.

Oostindjer et al. (2011a) also demonstrated a process of vertical social learning in which piglets learned to eat solid food from the sow, showing shorter latencies to eat, greater consumption and preference for the feed containing the same added flavour as was consumed by the sow and piglets during lactation. The present experiment failed to show any differences between treatments for investigative and feeding related behaviours in piglets, which did not support the hypotheses that these behaviours would occur more frequently in farrowing pens. However this result makes sense given that the feeding system in the farrowing pens did not allow piglets to actively observe the sow eating or to eat from the same container.

## **6. Conclusion**

In summary, the present experiment provided some evidence of improved welfare for sows and piglets housed in farrowing pens compared to farrowing crates during the lactation period. This was apparent from the increased interaction between sows and piglets and increased frequency of piglet play behavior and the decreased rate of manipulative behavior observed in farrowing pens. These behavioural differences were likely due to a combination of increased social and environmental enrichment provided by the larger space allowance and unrestrained sow in the farrowing pens. In contrast however, the increased sow injuries, displacements at the udder and piglets missing suckling bouts in farrowing pens presents some conflicting evidence with regard to welfare.

Whether the apparent welfare improvements of farrowing pens are significant enough to bring about long-term benefits for piglets and consequently improved sow productivity still requires further research. It would be beneficial to further explore the effect that sow experience has on sow behaviour in farrowing pens, which may confirm greater benefits of the housing system on sow and piglet productivity and welfare. Comparison of productivity measures between the two treatments was inconclusive and also requires further research. Importantly, comparison of productivity and welfare needs to follow piglets post-weaning, as some of the benefits of lactation housing may not become apparent until later in production.

## 7. Acknowledgements

Thank you to Geordie and Melville Charles and staff at Berrybank farm for your cooperation and assistance throughout the project.

Thank you also to the Australian Pork CRC for funding the project and to Professor Paul Hemsworth, Judy Nash and Maxine Rice for your guidance.

## References

Algers, B 1993, 'Nursing in pigs: communicating needs and distributing resources', *Journal of Animal Science*, vol. 71, no. 10, pp. 2826-31.

Algers, B & Uvnäs-Moberg, K 2007, 'Maternal behavior in pigs', *Hormones and Behavior*, vol. 52, pp. 78-85.

Andersen, IL, Berg, S & Bøe, KE 2005, 'Crushing of piglets by the mother sow (*Sus scrofa*)—purely accidental or a poor mother?', *Applied Animal Behaviour Science*, vol. 93, pp. 229-43.

Arey, DS & Sancha, ES 1996, 'Behaviour and productivity of sows and piglets in a family system and in farrowing crates', *Applied Animal Behaviour Science*, vol. 50, pp. 135-45.

Barnett, JL & Hemsworth, PH 2009, 'Welfare Monitoring Schemes: Using Research to Safeguard Welfare of Animals on the Farm', *Journal of Applied Animal Welfare Science*, vol. 12, pp. 114-31.

Barnett, JL, Hemsworth, PH, Cronin, GM, Jongman, EC & Hutson, GD 2001, 'A review of the welfare issues for sows and piglets in relation to housing', *Australian Journal of Agricultural Research*, vol. 52, no. 1, pp. 1-28.

Baxter, E, Jarvis, S, Sherwood, L, Farish, M, Roehe, R, Lawrence, AB & Edwards, SA 2011, 'Genetic and environmental effects on piglet survival and maternal behaviour of the farrowing sow', *Applied Animal Behaviour Science*, vol. 130, pp. 28-41.

Baxter, E, Lawrence, AB & Edwards, SA 2011a, 'Alternative farrowing accommodation: welfare and economic aspects of existing farrowing and lactation systems for pigs', *animal*, vol. 6, no. 1, pp. 96-117.

Baxter, EM, Lawrence, AB & Edwards, SA 2011b, 'Alternative farrowing systems: design criteria for farrowing systems based on the biological needs of sows and piglets', *animal*, vol. 5, no. 4, pp. 580-600.

Baxter, EM, Lawrence, AB & Edwards, SA 2012, 'Alternative farrowing accommodation: welfare and economic aspects of existing farrowing and lactation systems for pigs', *animal*, vol. 6, no. 01, pp. 96-117.

Baxter, MR 1991, 'The Freedom Farrowing System', *Farm Building Progress*, no. 104, pp. 9-15.

Blackshaw, JK, Blackshaw, AW, Thomas, FJ & Newman, FW 1994, 'Comparison of behaviour patterns of sows and litters in a farrowing crate and a farrowing pen', *Applied Animal Behaviour Science*, vol. 39, pp. 281-95.

Bøe, K 1993, 'Maternal behaviour of lactating sows in a loosehousing system', *Applied Animal Behaviour Science*, vol. 35, no. 4, pp. 327-38.

Boissy, A, Manteuffel, G, Jensen, MB, Moe, RO, Spruijt, B, Keeling, LJ, Winckler, C, Forkman, B, Dimitrov, I, Langbein, J, Bakken, M, Veissier, I & Aubert, A 2007, 'Assessment of positive emotions in animals to improve their welfare', *Physiology & Behavior*, vol. 92, no. 3, pp. 375-97.

Broom, DM 2008, 'Welfare Assessment and Relevant Ethical Decisions: Key Concepts', *ARBS Annual Review of Biomedical Sciences*, vol. 10, pp. T79-T90.

- Chaloupková, H, Illmann, G, Bartoš, L & Špinka, M 2007, 'The effect of pre-weaning housing on the play and agonistic behaviour of domestic pigs', *Applied Animal Behaviour Science*, vol. 103, no. 1-2, pp. 25-34.
- Council of Europe 2001, 'Commission Directive 2001/88/EC of 23 October 2001 amending Directive 91/630/EEC laying down minimum standards for the protection of pigs.', *Official Journal of the European communities*, vol. L316, pp. 1-3.
- Cronin, GM, Lef, bure, B & McClintock, S 2000, 'A comparison of piglet production and survival in the Werrabee Farrowing Pen and conventional farrowing crates at a commercial farm', *Australian Journal of Experimental Agriculture*, vol. 40, no. 1, pp. 17-23.
- Cronin, GM, Simpson, GJ & Hemsworth, PH 1996, 'The effects of the gestation and farrowing environments on sow and piglet behaviour and piglet survival and growth in early lactation', *Applied Animal Behaviour Science*, vol. 46, pp. 175-92.
- Cronin, GM & Smith, JA 1992, 'Suckling behaviour of sows in farrowing crates and straw-bedded pens', *Applied Animal Behaviour Science*, vol. 33, pp. 175-89.
- Dawkins, MS 1988, 'Behavioural deprivation: A central problem in animal welfare', *Applied Animal Behaviour Science*, vol. 20, no. 3-4, pp. 209-25.
- de Oliveira Junior, GM, Ferreira, AS, Oliviera, RFM, Silva, BAN, de Figueiredo, EM & Santos, M 2011, 'Behaviour and performance of lactating sows housed in different types of farrowing rooms during summer', *Livestock Science*, vol. 141, pp. 194-201.
- Dybkjær, L 1992, 'The identification of behavioural indicators of 'stress' in early weaned piglets', *Applied Animal Behaviour Science*, vol. 35, no. 2, pp. 135-47.
- Edwards, SA 2002, 'Perinatal mortality in the pig: environmental or physiological solutions?', *Livestock Production Science*, vol. 78, no. 1, pp. 3-12.
- Edwards, SA & Fraser, D 1997, 'Housing Systems for Farrowing and Lactation', *PIG JOURNAL*, vol. 39, pp. 77-89.
- Glastonbury, JRW 1976, 'A Survey of Prewearing Mortality in the Pig', *Australian Veterinary Journal*, vol. 52, no. 6, pp. 272-6.
- Gonyou, HW 2001, 'The Social Behaviour of Pigs', in LJ Keeling & HW Gonyou (eds), *Social Behaviour in Farm Animals*, CABI Publishing, Wallingford.
- Gu, Z, Gao, Y, Lin, B, Zhong, Z, Liu, Z, Wang, C & Li, B 2011, 'Impacts of a freedom farrowing pen design on sow behaviours and performance', *Preventive Veterinary Medicine*, vol. 102, pp. 296-303.
- Gustafsson, M, Jensen, P, de Jonge, FH, Illmann, G & Spinka, M 1999, 'Maternal behaviour of domestic sows and crosses between domestic sows and wild boar', *Applied Animal Behaviour Science*, vol. 65, no. 1, pp. 29-42.
- Held, SDE & Špinka, M 2011, 'Animal play and animal welfare', *Animal behaviour*, vol. 81, no. 5, pp. 891-9.
- Hemsworth, PH, Winfield, CG & Mullaney, PD 1976, 'A study of the development of the teat order in piglets', *Applied Animal Ethology*, vol. 2, no. 3, pp. 225-33.
- Horrell, I 1982, 'Social interactions amongst fostered piglets', *Applied Animal Ethology*, vol. 9, no. 1, pp. 85-6.
- Johnson, AK & Marchant-Forde, JN 2009, 'Welfare of Pigs in the Farrowing Environment', in JN Marchant-Forde (ed.), *The Welfare of Pigs*, Springer Science and Business Media, pp. 141-88.
- Karlen, GAM, Hemsworth, PH, Gonyou, HW, Fabrega, E, Strom, AD & Smits, RJ 2007, 'The welfare of gestating sows in conventional stalls and large groups on deep litter', *Applied Animal Behaviour Science*, vol. 105, pp. 87-101.
- KilBride, AL, Mendl, M, Statham, P, Held, S, Harris, M, Cooper, S & Green, LE 2012, 'A cohort study of preweaning piglet mortality and farrowing accommodation on 112 commercial pig farms in England', *Preventive Veterinary Medicine*, vol. 104, no. 3-4, pp. 281-91.

- Marchant, JN, Rudd, AR, Mendl, MT, Broom, DM, Meredith, MJ, Corning, S & Simmins, PH 2000, 'Timing and causes of piglet mortality in alternative and conventional farrowing systems', *Veterinary Record*, vol. 147, no. 8, pp. 209-14.
- Morrow-Tesch, J & McGlone, JJ 1990a, 'Sensory systems and nipple attachment behavior in neonatal pigs', *Physiology & Behavior*, vol. 47, no. 1, pp. 1-4.
- Morrow-Tesch, J & McGlone, JJ 1990b, 'Sources of maternal odors and the development of odor preferences in baby pigs', *Journal of Animal Science*, vol. 68, no. 11, pp. 3563-71.
- Newberry, RC, Wood-Gush, DGM & Hall, JW 1988, 'Playful behaviour of piglets', *Behavioural Processes*, vol. 17, no. 3, pp. 205-16.
- Oostindjer, M, Bolhuis, JE, Mendl, M, Held, S, Gerrits, W, van den Brand, H & Kemp, B 2010, 'Effects of environmental enrichment and loose housing of lactating sows on piglet performance before and after weaning', *Journal of Animal Science*, vol. 88, no. 11, pp. 3554-62.
- Oostindjer, M, Bolhuis, JE, Mendl, M, Held, S, van den Brand, H & Kemp, B 2011a, 'Learning how to eat like a pig: effectiveness of mechanisms for vertical social learning in piglets', *Animal behaviour*, vol. 82, no. 3, pp. 503-11.
- Oostindjer, M, van den Brand, H, Kemp, B & Bolhuis, J 2011b, 'Effects of environmental enrichment and loose housing of lactating sows on piglet behaviour before and after weaning', *Applied Animal Behaviour Science*, vol. 134, no. 1-2, pp. 31-41.
- Oostindjer, M, van den Brand, H, Kemp, B & Bolhuis, JE 2011c, 'Effects of environmental enrichment and loose housing of lactating sows on piglet behaviour before and after weaning', *Applied Animal Behaviour Science*, vol. 134, pp. 31-41.
- Pedersen, AR, Moustsen, VA, Nielsen, MBF & Kristensen, AR 2011, 'Improved udder access prolongs duration of milk letdown and increases piglet weight gain', *Livestock Science*, vol. 140, pp. 253-61.
- Pedersen, LJ, Jørgensen, E, Heiskanen, T & Damm, BI 2006, 'Early piglet mortality in loose-housed sows related to sow and piglet behaviour and to the progress of parturition', *Applied Animal Behaviour Science*, vol. 96, no. 3-4, pp. 215-32.
- PISC 2008, *Model code of practice for the welfare of animals: pigs*, 3rd ed. PISC Report 92, CSIRO Publishing, Victoria, Australia.
- Straw, BE & Bartlett, P 2001, 'Flank or belly nosing in weaned pigs', *Journal of Swine Health and Production*, vol. 9, no. 1, pp. 19-23.
- Weber, R, Keil, NM, Fehr, M & Horat, R 2009, 'Factors affecting piglet mortality in loose farrowing systems on commercial farms', *Livestock Science*, vol. 124, no. 1-3, pp. 216-22.
- Wechsler, B 1996, 'Rearing pigs in species-specific family groups', *Animal Welfare*, vol. 5, no. 1, pp. 25-35.